

AN EXTENSION OF A CONGRUENCE BY TAURASO

ROMEO MEŠTROVIĆ

ABSTRACT. For a positive integer n let $H_n = \sum_{k=1}^n 1/k$ be the n th harmonic number. In this note we prove that for any prime $p \geq 7$,

$$\sum_{k=1}^{p-1} \frac{H_k}{k^2} \equiv \sum_{k=1}^{p-1} \frac{H_k^2}{k} \equiv \frac{3}{2p} \sum_{k=1}^{p-1} \frac{1}{k^2} \pmod{p^2}.$$

Notice that the first part of this congruence is recently proposed by R. Tauraso as a problem in Amer. Math. Monthly. In our elementary proof of the second part of the above congruence we use certain classical congruences modulo a prime and the square of a prime, some congruences involving harmonic numbers and a combinatorial identity due to V. Hernández.

1. INTRODUCTION AND MAIN RESULTS

Given positive integers n and m , the *harmonic numbers of order m* are those rational numbers $H_{n,m}$ defined as

$$H_{n,m} = \sum_{k=1}^n \frac{1}{k^m}.$$

For simplicity, we will denote by

$$H_n := H_{n,1} = \sum_{k=1}^n \frac{1}{k}$$

the n th *harmonic number* (we assume in addition that $H_{0,m} = H_0 = 0$).

Usually, here as always in the sequel, we consider the congruence relation modulo a prime p extended to the ring of rational numbers with denominators not divisible by p . For such fractions we put $m/n \equiv r/s \pmod{p}$ if and only if $ms \equiv nr \pmod{p}$, and the residue class of m/n is the residue class of mn' where n' is the inverse of n modulo p .

By a problem proposed by R. Tauraso in [12] and recently solved by D. B. Tyler [15], for any prime $p \geq 7$,

$$(1) \quad \sum_{k=1}^{p-1} \frac{H_k}{k^2} \equiv \sum_{k=1}^{p-1} \frac{H_k^2}{k} \pmod{p^2}.$$

2010 *Mathematics Subject Classification*. Primary 11B75, 11A07; Secondary 11B65, 05A19, 05A10.

Keywords and phrases. congruence modulo prime (prime power), n th harmonic number of order m , Bernoulli number

Further, R. Tauraso [14, Theorem 2.3] proved

$$(2) \quad \sum_{k=1}^{p-1} \frac{H_k}{k^2} \equiv -\frac{3}{p^2} \sum_{k=1}^{p-1} \frac{1}{k} \pmod{p^2}.$$

Tauraso's proof of (2) is based on an identity due to V. Hernández [4] (see Lemma 2.5) and the congruence for triple harmonic sum modulo a prime due to Zhao [16] (see (47) of Remarks in Section 2). In this note, we give an elementary proof of (2) and its extension as follows.

Theorem 1.1. *If $p \geq 7$ is a prime, then*

$$(3) \quad \sum_{k=1}^{p-1} \frac{H_k}{k^2} \equiv \sum_{k=1}^{p-1} \frac{H_k^2}{k} \equiv -\frac{3}{p^2} \sum_{k=1}^{p-1} \frac{1}{k} \equiv \frac{3}{2p} \sum_{k=1}^{p-1} \frac{1}{k^2} \pmod{p^2}.$$

Recall that Z. W. Sun in [10] established basic congruences modulo a prime $p \geq 5$ for several sums of terms involving harmonic numbers. In particular, Sun established $\sum_{k=1}^{p-1} H_k^r \pmod{p^{4-r}}$ for $r = 1, 2, 3$. Further generalizations of these congruences are recently obtained by Tauraso in [13].

Recall that the *Bernoulli numbers* B_k are defined by the generating function

$$\sum_{k=0}^{\infty} B_k \frac{x^k}{k!} = \frac{x}{e^x - 1}.$$

It is easy to find the values $B_0 = 1$, $B_1 = -\frac{1}{2}$, $B_2 = \frac{1}{6}$, $B_4 = -\frac{1}{30}$, and $B_n = 0$ for odd $n \geq 3$. Furthermore, $(-1)^{n-1} B_{2n} > 0$ for all $n \geq 1$. Applying a congruence given in [9, Theorem 5.1(a)] related to the sum $\sum_{k=1}^{p-1} 1/k^2$ modulo p^3 , the congruence (3) in terms of Bernoulli numbers may be written as follows.

Corollary 1.2. *Let $p \geq 7$ be a prime. Then*

$$(4) \quad \sum_{k=1}^{p-1} \frac{H_k^2}{k} \equiv \sum_{k=1}^{p-1} \frac{H_k}{k^2} \equiv 3 \left(\frac{B_{2p-4}}{2p-4} - \frac{2B_{p-3}}{p-3} \right) \pmod{p^2}.$$

In particular, we have

$$(5) \quad \sum_{k=1}^{p-1} \frac{H_k^2}{k} \equiv \sum_{k=1}^{p-1} \frac{H_k}{k^2} \equiv B_{p-3} \pmod{p}.$$

Remark. Notice that the second congruence of (5) was obtained by Sun and Tauraso [11, the congruence (5.4)] by using a standard technique expressing sum of powers in terms of Bernoulli numbers.

Our proof of the second part of the congruence (3) given in the next section is entirely elementary. It is based on certain classical congruences modulo a prime and the square of a prime, two simple congruences given by Z. W. Sun [10] and two particular cases of a combinatorial identity due to V. Hernández [4].

2. PROOF OF THEOREM 1.1

The following congruences by Z. W. Sun given in his recent paper [10] are needed in the proof of Theorem 1.1.

Lemma 2.1. *Let $p \geq 7$ be a prime. Then*

$$(6) \quad H_{p-k} \equiv H_{k-1} \pmod{p}$$

and

$$(7) \quad (-1)^k \binom{p-1}{k} \equiv 1 - pH_k + \frac{p^2}{2}(H_k^2 - H_{k,2}) \pmod{p^3}$$

for every $k = 1, 2, \dots, p-1$.

Proof. The congruences (6) and (7) are in fact the congruences (2.1) and (2.2) in [10, Lemma 2.1], respectively. $\square$

The following well known result is a generalization of *Wolstenholme's theorem* (see, e.g., [1, Theorem 1] or [3]).

Lemma 2.2. ([2, Theorem 3]). *Let m be a positive integer and let p be a prime such that $p \geq m+3$. Then*

$$(8) \quad H_{p-1,m} \equiv \begin{cases} 0 & \pmod{p} \text{ if } m \text{ is even} \\ 0 & \pmod{p^2} \text{ if } m \text{ is odd.} \end{cases}$$

In particular, for any prime $p \geq 5$,

$$(9) \quad H_{p-1} \equiv 0 \pmod{p^2} \quad (\text{Wolstenholme's theorem}),$$

and for any prime $p \geq 7$, $H_{p-1,3} \equiv 0 \pmod{p^2}$ and $H_{p-1,2} \equiv H_{p-1,4} \equiv 0 \pmod{p}$.

Lemma 2.3. *Let $p \geq 7$ be a prime. Then*

$$(10) \quad \sum_{k=1}^n \frac{H_{k-1}}{k^3} \equiv \sum_{k=1}^n \frac{H_k}{k^3} \equiv 0 \pmod{p},$$

$$(11) \quad \sum_{k=1}^{p-1} \frac{H_k^3}{k} \equiv \frac{3}{2} \sum_{k=1}^{p-1} \frac{H_k^2}{k^2} \pmod{p},$$

$$(12) \quad \sum_{k=1}^n \frac{H_{k-1,3}}{k} \equiv \sum_{k=1}^n \frac{H_{k,3}}{k} \equiv 0 \pmod{p},$$

and

$$(13) \quad \sum_{k=1}^{p-1} \frac{H_k}{k^2} \equiv \sum_{k=1}^{p-1} \frac{H_k^2}{k} \pmod{p^2}.$$

Proof. By the congruence (6) from Lemma 2.1, $H_k \equiv H_{p-k-1} \pmod{p}$ for each $k = 1, 2, \dots, p-1$ (notice that this is true for $k = p-1$ because $p \mid H_{p-1}$), and therefore

$$(14) \quad \sum_{k=1}^{p-1} \frac{H_{k-1}}{k^3} = \sum_{k=1}^{p-1} \frac{H_{(p-k)-1}}{(p-k)^3} \equiv \sum_{k=1}^{p-1} \frac{H_k}{(p-k)^3} \equiv - \sum_{k=1}^{p-1} \frac{H_k}{k^3} \pmod{p}.$$

Furthermore, using (8) with $m = 4$ we get

$$(15) \quad \sum_{k=1}^{p-1} \frac{H_{k-1}}{k^3} = \sum_{k=1}^{p-1} \frac{(H_k - \frac{1}{k})}{k^3} = \sum_{k=1}^{p-1} \frac{H_k}{k^3} - \sum_{k=1}^{p-1} \frac{1}{k^4} \equiv \sum_{k=1}^{p-1} \frac{H_k}{k^3} \pmod{p}.$$

From (14) and (15) it follows that

$$\sum_{k=1}^{p-1} \frac{H_{k-1}}{k^3} \equiv \sum_{k=1}^{p-1} \frac{H_k}{k^3} \equiv 0 \pmod{p},$$

which is actually (10).

Since $H_k = H_{k-1} + 1/k$, for each $k = 1, 2, \dots, p-1$ we have

$$\begin{aligned} \frac{H_k^3}{k} - \frac{H_{k-1}^3}{k} &= \frac{1}{k}(H_k - H_{k-1}) \left(H_k^2 + H_k \left(H_k - \frac{1}{k} \right) + \left(H_k - \frac{1}{k} \right)^2 \right) \\ &= \frac{1}{k^2} \left(3H_k^2 - 3\frac{H_k}{k} + \frac{1}{k^2} \right) = 3\frac{H_k^2}{k^2} - 3\frac{H_k}{k^3} + \frac{1}{k^4}. \end{aligned}$$

The above identity, (10) and (8) of Lemma 2.2 with $m = 4$ yield

$$(16) \quad \sum_{k=1}^{p-1} \frac{H_k^3}{k} - \sum_{k=1}^{p-1} \frac{H_{k-1}^3}{k} = 3 \sum_{k=1}^{p-1} \frac{H_k^2}{k^2} - 3 \sum_{k=1}^{p-1} \frac{H_k}{k^3} + \sum_{k=1}^{p-1} \frac{1}{k^4} \equiv 3 \sum_{k=1}^{p-1} \frac{H_k^2}{k^2} \pmod{p}.$$

On the other hand, since by (6) from Lemma 2.1, $H_k \equiv H_{p-k-1} \pmod{p}$ for each $k = 1, 2, \dots, p-1$, then

$$(17) \quad \sum_{k=1}^{p-1} \frac{H_{k-1}^3}{k} = \sum_{k=1}^{p-1} \frac{H_{(p-k)-1}^3}{p-k} \equiv \sum_{k=1}^{p-1} \frac{H_k^3}{p-k} \equiv - \sum_{k=1}^{p-1} \frac{H_k^3}{k} \pmod{p}.$$

Taking (17) into (16) gives

$$(18) \quad \sum_{k=1}^{p-1} \frac{H_k^3}{k} \equiv \frac{3}{2} \sum_{k=1}^{p-1} \frac{H_k^2}{k^2} \pmod{p},$$

which proves (11).

Proof of the congruence (12) is completely analogous to the previous proof using the fact that by Lemma 2.2, $H_{p-1,3} \equiv 0 \pmod{p}$ for each $k = 1, \dots, p-1$, and therefore

$$H_{k-1,3} = \sum_{i=1}^{k-1} \frac{1}{i^3} = H_{p-1,3} - \sum_{j=1}^{p-k} \frac{1}{(p-j)^3} \equiv \sum_{j=1}^{p-k} \frac{1}{j^3} = H_{p-k,3} \pmod{p}.$$

Finally (cf. [15]), from the identity

$$H_{k-1}^3 = \left(H_k - \frac{1}{k}\right)^3 = H_k^3 - 3\frac{H_k^2}{k} + 3\frac{H_k}{k^2} - \frac{1}{k^3}$$

immediately follows that

$$(19) \quad \sum_{k=1}^{p-1} \frac{H_k^2}{k} - \sum_{k=1}^{p-1} \frac{H_k}{k^2} = \frac{1}{3} (H_{p-1}^3 - H_{p-1,3}).$$

Inserting in the right hand side of the identity (19) the congruences $H_{p-1} \equiv H_{p-1,3} \equiv 0 \pmod{p^2}$ given in Lemma 2.2, we immediately obtain (13). This completes the proof. $\square$

Lemma 2.4. *Let $p \geq 7$ be a prime. Then*

$$(20) \quad \sum_{k=1}^{p-1} \frac{H_k \cdot H_{k,2}}{k} \equiv \sum_{1 \leq i < j < k \leq p-1} \frac{1}{i^2 j k} + \sum_{1 \leq i < j < k \leq p-1} \frac{1}{i^2 j k} \pmod{p}.$$

Proof. Since $H_k = H_{k-1} + 1/k$ for every $k = 1, 2, \dots, p-1$, we get

$$(21) \quad \begin{aligned} \sum_{k=1}^{p-1} \frac{H_k \cdot H_{k,2}}{k} &= \sum_{k=1}^{p-1} \frac{1}{k} \left(H_{k-1} + \frac{1}{k}\right) \left(H_{k-1,2} + \frac{1}{k^2}\right) \\ &= \sum_{k=1}^{p-1} \frac{H_{k-1} \cdot H_{k-1,2}}{k} + \sum_{k=1}^{p-1} \frac{H_{k-1,2}}{k^2} + \sum_{k=1}^{p-1} \frac{H_{k-1}}{k^3} + \sum_{k=1}^{p-1} \frac{1}{k^4}. \end{aligned}$$

Using particular congruences given in Lemma 2.2 with $m = 2$ and $m = 4$, we find that

$$(22) \quad \begin{aligned} \sum_{k=1}^{p-1} \frac{H_{k-1,2}}{k^2} &= \sum_{k=1}^{p-1} \frac{1}{k^2} \sum_{i=1}^{k-1} \frac{1}{i^2} = \sum_{1 \leq i < k \leq p-1} \frac{1}{i^2 k^2} \\ &= \frac{1}{2} \left(\sum_{k=1}^{p-1} \frac{1}{k^2} \right)^2 - \frac{1}{2} \sum_{k=1}^{p-1} \frac{1}{k^4} \equiv 0 \pmod{p}. \end{aligned}$$

Substituting the congruences (22), (10) of Lemma 2.3 and (8) with $m = 4$ of Lemma 2.2 into (21), we obtain

$$(23) \quad \sum_{k=1}^{p-1} \frac{H_k \cdot H_{k,2}}{k} \equiv \sum_{k=1}^{p-1} \frac{H_{k-1} \cdot H_{k-1,2}}{k} \pmod{p}.$$

The right hand side of (23) can be expressed as

$$\begin{aligned}
 (24) \quad \sum_{k=1}^{p-1} \frac{H_{k-1} \cdot H_{k-1,2}}{k} &= \sum_{k=1}^{p-1} \frac{1}{k} \left(1 + \frac{1}{2} + \cdots + \frac{1}{k-1} \right) \left(1 + \frac{1}{2^2} + \cdots + \frac{1}{(k-1)^2} \right) \\
 &= \sum_{k=1}^{p-1} \frac{1}{k} \left(\sum_{1 \leq i < j \leq k-1} \frac{1}{ij^2} + \sum_{1 \leq i < j \leq k-1} \frac{1}{i^2 j} + \sum_{i=1}^{k-1} \frac{1}{i^3} \right) \\
 &= \sum_{1 \leq i < j < k \leq p-1} \frac{1}{ij^2 k} + \sum_{1 \leq i < j < k \leq p-1} \frac{1}{i^2 j k} + \sum_{k=1}^{p-1} \frac{H_{k-1,3}}{k}.
 \end{aligned}$$

Taking (12) of Lemma 2.3 into (24), and comparing this with (23), we immediately obtain (20). $\square$

Further, for the proof of Theorem 1.1 we will need two particular cases of the following identity due to V. Hernández [4].

Lemma 2.5. ([4]) *Let n and m be positive integers. Then*

$$(25) \quad \sum_{k=1}^n \binom{n}{k} (-1)^{k-1} \sum_{1 \leq i_1 \leq i_2 \leq \cdots \leq i_m = k} \frac{1}{i_1 i_2 \cdots i_m} = \sum_{k=1}^n \frac{1}{k^m}.$$

Lemma 2.6. *Let $p \geq 7$ be a prime. Then*

$$(26) \quad \sum_{k=1}^{p-1} \frac{H_k \cdot H_{k,2}}{k} \equiv -\frac{3}{2} \sum_{k=1}^{p-1} \frac{H_k^2}{k^2} \pmod{p}.$$

Proof. The identity (25) of Lemma 2.5 with $m = 3$ and $n = p - 1$ becomes

$$(27) \quad \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} \binom{p-1}{k} \sum_{1 \leq i \leq j \leq k} \frac{1}{ij} = \sum_{k=1}^{p-1} \frac{1}{k^3} = H_{p-1,3}.$$

For any fixed $k \leq p - 1$, we have the identity

$$(28) \quad \sum_{1 \leq i \leq j \leq k} \frac{1}{ij} = \frac{1}{2} \left(\left(\sum_{i=1}^k \frac{1}{i} \right)^2 + \sum_{i=1}^k \frac{1}{i^2} \right) = \frac{1}{2} (H_k^2 + H_{k,2}).$$

Next the congruence (7) from Lemma 2.1 reduced modulo p^2 gives

$$(29) \quad (-1)^{k-1} \binom{p-1}{k} \equiv pH_k - 1 \pmod{p^2}$$

for every $k = 1, 2, \dots, p - 1$. Substituting (28), (29) and the congruence $H_{p-1,3} \equiv 0 \pmod{p^2}$ of Lemma 2.2 into (27), we immediately obtain

$$\sum_{k=1}^{p-1} \frac{1}{k} (pH_k - 1) (H_k^2 + H_{k,2}) \equiv 0 \pmod{p^2},$$

or equivalently,

$$(30) \quad p \left(\sum_{k=1}^{p-1} \frac{H_k^3}{k} + \sum_{k=1}^{p-1} \frac{H_k \cdot H_{k,2}}{k} \right) - \left(\sum_{k=1}^{p-1} \frac{H_k^2}{k} + \sum_{k=1}^{p-1} \frac{H_{k,2}}{k} \right) \equiv 0 \pmod{p^2}.$$

Further, (13) from Lemma 2.3 and the congruences $H_{p-1} \equiv H_{p-1,3} \equiv 0 \pmod{p^2}$ from Lemma 2.2 give

$$(31) \quad \begin{aligned} \sum_{k=1}^{p-1} \frac{H_k^2}{k} + \sum_{k=1}^{p-1} \frac{H_{k,2}}{k} &\equiv \sum_{k=1}^{p-1} \frac{H_k}{k^2} + \sum_{k=1}^{p-1} \frac{H_{k,2}}{k} \pmod{p^2} \\ &= \sum_{1 \leq i \leq k \leq p-1} \frac{1}{ik^2} + \sum_{1 \leq i \leq k \leq p-1} \frac{1}{i^2k} \\ &= \left(\sum_{i=1}^{p-1} \frac{1}{i} \right) \left(\sum_{k=1}^{p-1} \frac{1}{k^2} \right) + \sum_{k=1}^{p-1} \frac{1}{i^3} \\ &= H_{p-1} \cdot H_{p-1,2} + H_{p-1,3} \equiv 0 \pmod{p^2}. \end{aligned}$$

Substituting (31) into (30), we find that

$$(32) \quad \sum_{k=1}^{p-1} \frac{H_k^3}{k} + \sum_{k=1}^{p-1} \frac{H_k \cdot H_{k,2}}{k} \equiv 0 \pmod{p}.$$

Taking (11) of Lemma 2.3 into (32) yields (26). This concludes the proof. $\square$

Lemma 2.7. *Let $p \geq 7$ be a prime. Then*

$$(33) \quad \sum_{1 \leq i < j < k \leq p-1} \frac{1}{i^2jk} \equiv \sum_{1 \leq i < j < k \leq p-1} \frac{1}{ijk^2} \equiv -\frac{1}{2} \sum_{1 \leq i < j < k \leq p-1} \frac{1}{ij^2k} \pmod{p}.$$

Proof. For simplicity, we denote

$$A := \sum_{1 \leq i < j < k \leq p-1} \frac{1}{i^2jk}, \quad B := \sum_{1 \leq i < j < k \leq p-1} \frac{1}{ij^2k}, \quad \text{and} \quad C := \sum_{1 \leq i < j < k \leq p-1} \frac{1}{ijk^2}.$$

Obviously, holds the identity

$$(34) \quad \left(\sum_{1 \leq i < j < k \leq p-1} \frac{1}{ijk} \right) \left(\sum_{l=1}^{p-1} \frac{1}{l} \right) = A + B + C + 4 \sum_{1 \leq i < j < k < l \leq p-1} \frac{1}{ijkl}.$$

The well known Newton's identities (see e.g., [7]) imply

$$\begin{aligned} 4 \sum_{1 \leq i < j < k < l \leq p-1} \frac{1}{ijkl} &= \sum_{1 \leq i < j < k \leq p-1} \frac{1}{ijk} H_{p-1} - \sum_{1 \leq i < j \leq p-1} \frac{1}{ij} H_{p-1,2} \\ &\quad + \sum_{i=1}^{p-1} \frac{1}{i} H_{p-1,3} - H_{p-1,4}, \end{aligned}$$

whence since all the sums H_{p-1} , $H_{p-1,2}$, $H_{p-1,3}$ and $H_{p-1,4}$ are divisible by a prime $p \geq 7$, we obtain (cf. [16, Theorem 1.5] or [17])

$$(35) \quad \sum_{1 \leq i < j < k < l \leq p-1} \frac{1}{ijkl} \equiv 0 \pmod{p}.$$

Inserting (35) and $H_{p-1} = \sum_{i=1}^{p-1} 1/i \equiv 0 \pmod{p}$ into (34), we get

$$(36) \quad A + B + C \equiv 0 \pmod{p}.$$

Further, by the substitution trick $i, j, k \rightarrow p-i, p-j, p-k$,

$$(37) \quad \begin{aligned} A &= \sum_{1 \leq i < j < k \leq p-1} \frac{1}{i^2 j k} = \sum_{1 \leq p-i < p-j < p-k \leq p-1} \frac{1}{(p-i)^2 (p-j)(p-k)} \\ &\equiv \sum_{1 \leq k < j < i \leq p-1} \frac{1}{k j i^2} = C \pmod{p}. \end{aligned}$$

From (37) we see that $C \equiv A \pmod{p}$, which substituting into (36) gives

$$(38) \quad 2A + B \equiv 0 \pmod{p}.$$

Finally, (37) and (38) yield $C \equiv A \equiv -B/2 \pmod{p}$, as desired. $\square$

Lemma 2.8. *Let $p \geq 7$ be a prime. Then*

$$(39) \quad \sum_{k=1}^{p-1} \frac{H_k^2}{k^2} \equiv - \sum_{1 \leq i < j < k \leq p-1} \frac{1}{ij^2 k} \pmod{p}.$$

Proof. We follow proof of the congruence (3) in Theorem 1.1 of [5]. By Lemma 2.2, $H_{p-1} := \sum_{j=1}^{p-1} 1/j \equiv 0 \pmod{p}$, or equivalently, for each $j = 1, 2, \dots, p-2$ holds

$$(40) \quad \frac{1}{j+1} + \frac{1}{j+2} + \dots + \frac{1}{p-1} \equiv - \left(1 + \frac{1}{2} + \dots + \frac{1}{j-1} + \frac{1}{j} \right) \pmod{p^2}.$$

Applying the congruence (40), we find that

$$\begin{aligned}
\sum_{1 \leq i < j < k \leq p-1} \frac{1}{ij^2k} &= \sum_{j=2}^{p-1} \frac{1}{j^2} \sum_{i=1}^{j-1} \frac{1}{i} \sum_{k=j+1}^{p-1} \frac{1}{k} \\
&= \sum_{j=2}^{p-1} \frac{1}{j^2} \left(1 + \frac{1}{2} + \cdots + \frac{1}{j-1}\right) \left(\frac{1}{j+1} + \frac{1}{j+2} + \cdots + \frac{1}{p-1}\right) \\
&\equiv \sum_{j=2}^{p-1} \frac{1}{j^2} \left(1 + \frac{1}{2} + \cdots + \frac{1}{j-1}\right) \left(-\left(1 + \frac{1}{2} + \cdots + \frac{1}{j}\right)\right) \pmod{p} \\
&= \sum_{j=1}^{p-1} \frac{1}{j^2} \left(H_j - \frac{1}{j}\right) (-H_j) = -\sum_{j=1}^{p-1} \frac{H_j^2}{j^2} + \sum_{j=1}^{p-1} \frac{H_j}{j^3} \\
&= -\sum_{j=1}^{p-1} \frac{H_j^2}{j^2} + \sum_{j=1}^{p-1} \frac{H_{j-1} + \frac{1}{j}}{j^3} = -\sum_{j=1}^{p-1} \frac{H_j^2}{j^2} + \sum_{1 \leq i < j \leq p-1} \frac{1}{ij^3} + \sum_{j=1}^{p-1} \frac{1}{j^4} \\
&= -\sum_{j=1}^{p-1} \frac{H_j^2}{j^2} + \sum_{j=1}^{p-1} \frac{H_{j-1}}{j^3} + H_{p-1,4} \pmod{p},
\end{aligned}$$

whence it follows that

$$(41) \quad \sum_{j=1}^{p-1} \frac{H_j^2}{j^2} \equiv \sum_{j=1}^{p-1} \frac{H_{j-1}}{j^3} + H_{p-1,4} - \sum_{1 \leq i < j < k \leq p-1} \frac{1}{ij^2k} \pmod{p}.$$

Since by the first part of (10) from Lemma 2.3 and by (8) of Lemma 2.2 with $m = 4$,

$$\sum_{j=1}^{p-1} \frac{H_{j-1}}{j^3} \equiv H_{p-1,4} \equiv 0 \pmod{p},$$

substituting this into (41), we obtain (39). $\square$

The first congruence of the following result was recently established by Z. W. Sun [10, Theorem 1.1 (1.5)].

Lemma 2.9. *Let $p \geq 7$ be a prime. Then*

$$(42) \quad \sum_{k=1}^{p-1} \frac{H_k^2}{k^2} \equiv 0 \pmod{p},$$

$$(43) \quad \sum_{k=1}^{p-1} \frac{H_k \cdot H_{k,2}}{k} \equiv 0 \pmod{p}$$

and

$$(44) \quad \sum_{k=1}^{p-1} \frac{H_k^3}{k} \equiv 0 \pmod{p}.$$

Proof. Comparing the congruences (20) of Lemma 2.4 and (26) of Lemma 2.6, we have

$$(45) \quad \sum_{k=1}^{p-1} \frac{H_k^2}{k^2} \equiv -\frac{2}{3} \left(\sum_{1 \leq i < j < k \leq p-1} \frac{1}{ij^2k} + \sum_{1 \leq i < j < k \leq p-1} \frac{1}{i^2jk} \right) \pmod{p}.$$

Since by (33) of Lemma 2.7,

$$\sum_{1 \leq i < j < k \leq p-1} \frac{1}{i^2jk} \equiv -\frac{1}{2} \sum_{1 \leq i < j < k \leq p-1} \frac{1}{ij^2k} \pmod{p},$$

then substituting this into (45), we obtain

$$(46) \quad \sum_{k=1}^{p-1} \frac{H_k^2}{k^2} \equiv -\frac{1}{3} \sum_{1 \leq i < j < k \leq p-1} \frac{1}{ij^2k} \pmod{p}.$$

Finally, as by (39) of Lemma 2.8,

$$\sum_{k=1}^{p-1} \frac{H_k^2}{k^2} \equiv - \sum_{1 \leq i < j < k \leq p-1} \frac{1}{ij^2k} \pmod{p},$$

then comparing this with (46) implies

$$\sum_{k=1}^{p-1} \frac{H_k^2}{k^2} \equiv 0 \pmod{p},$$

which coincides with (42).

Finally, (42) and (26) of Lemma 2.6 yield (43), while (42) and (11) of Lemma 2.3 yield (44). $\square$

Remarks. Applying a standard technique expressing sum of powers in terms of Bernoulli numbers, Z. W. Sun in [10, Proof of (1.5) of Theorem] showed that

$$\sum_{k=1}^{p-1} \frac{H_k^2}{k^2} \equiv - \sum_{j=0}^{p-3} B_j B_{p-3-j} \pmod{p}.$$

The above congruence and (42) yield the following curious congruence for a prime $p \geq 7$ established by J. Zhao [16, (3.19) of Corollary 3.6]:

$$\sum_{j=0}^{p-3} B_j B_{p-3-j} \equiv 0 \pmod{p}.$$

As noticed in [10, Proof of Lemma 2.8], the above congruence immediately follows from an identity of Matiyasevich (cf. [8, (1.3)]).

Furthermore, the congruences (42), (46) and (33) from Lemma 2.7 immediately give

$$(47) \quad \sum_{1 \leq i < j < k \leq p-1} \frac{1}{i^2jk} \equiv \sum_{1 \leq i < j < k \leq p-1} \frac{1}{ij^2k} \equiv \sum_{1 \leq i < j < k \leq p-1} \frac{1}{ij^2k} \equiv 0 \pmod{p}.$$

Notice that the congruences (47) were proved by J. Zhao [16, Corollary 3.6 (3.20)] applying a technique expressing sum of powers in terms of Bernoulli numbers.

The following result is contained in [6, Lemma 2.4].

Lemma 2.10. *Let $p \geq 7$ be a prime. Then*

$$(48) \quad 2 \sum_{k=1}^{p-1} \frac{1}{k} \equiv -p \sum_{k=1}^{p-1} \frac{1}{k^2} \pmod{p^4}.$$

Proof. Multiplying the identity

$$1 + \frac{p}{k} + \frac{p^2}{k^2} = \frac{p^3 - k^3}{k^2(p - k)}$$

by $-p/k^2$ ($1 \leq k \leq p-1$), we obtain

$$-\frac{p}{k^2} \left(1 + \frac{p}{k} + \frac{p^2}{k^2} \right) = \frac{-p^4 + pk^3}{k^4(p - k)} \equiv \frac{p}{k(p - k)} \pmod{p^4},$$

which can be written as

$$\frac{1}{k} + \frac{1}{p - k} \equiv - \left(\frac{p}{k^2} + \frac{p^2}{k^3} + \frac{p^3}{k^4} \right) \pmod{p^4}.$$

After summation of the above congruence over $k = 1, \dots, p-1$ we immediately obtain

$$(49) \quad 2H_{p-1} \equiv -pH_{p-1,2} - p^2H_{p-1,3} - p^3H_{p-1,4} \pmod{p^4}.$$

Taking the congruences $H_{p-1,3} \equiv 0 \pmod{p^2}$ and $H_{p-1,4} \equiv 0 \pmod{p}$ of Lemma 2.2 into (49), it becomes (48). $\square$

Proof of Theorem 1.1. The first congruence of (3) is in fact the congruence (13) of Lemma 2.3.

The identity (25) of Lemma 2.5 with $m = 2$ and $n = p-1$ becomes

$$(50) \quad \sum_{1 \leq i \leq k \leq p-1} (-1)^{k-1} \binom{p-1}{k} \frac{1}{ik} = \sum_{k=1}^{p-1} \frac{1}{k^2} = H_{p-1,2}.$$

Substituting the congruence (7) of Lemma 2.1 into the left hand side of equality (50), we find that

$$(51) \quad \sum_{1 \leq i \leq k \leq p-1} \frac{(-1)^{k-1}}{ik} \binom{p-1}{k} \equiv - \sum_{1 \leq i \leq k \leq p-1} \frac{1}{ik} + p \sum_{1 \leq i \leq k \leq p-1} \frac{H_k}{ik} - \frac{p^2}{2} \sum_{1 \leq i \leq k \leq p-1} \frac{H_k^2}{ik} + \frac{p^2}{2} \sum_{1 \leq i \leq k \leq p-1} \frac{H_{k,2}}{ik} \pmod{p^3}.$$

By Wolstenholme's theorem,

$$(52) \quad \sum_{1 \leq i \leq k \leq p-1} \frac{1}{ik} = \frac{1}{2} (H_{p-1}^2 + H_{p-1,2}) \equiv \frac{1}{2} H_{p-1,2} \pmod{p^2}.$$

Next we have

$$(53) \quad \sum_{1 \leq i \leq k \leq p-1} \frac{H_k}{ik} = \sum_{k=1}^{p-1} \frac{H_k}{k} \sum_{i=1}^k \frac{1}{i} = \sum_{k=1}^{p-1} \frac{H_k^2}{k}.$$

Further, using (44) of Lemma 2.9,

$$(54) \quad \sum_{1 \leq i \leq k \leq p-1} \frac{H_k^2}{ik} = \sum_{k=1}^{p-1} \frac{H_k^2}{k} \sum_{i=1}^k \frac{1}{i} = \sum_{k=1}^{p-1} \frac{H_k^3}{k} \equiv 0 \pmod{p}.$$

Similarly, using (43) of Lemma 2.9,

$$(55) \quad \sum_{1 \leq i \leq k \leq p-1} \frac{H_{k,2}}{ik} = \sum_{k=1}^{p-1} \frac{H_{k,2}}{k} \sum_{i=1}^k \frac{1}{i} = \sum_{k=1}^{p-1} \frac{H_{k,2} \cdot H_k}{k} \equiv 0 \pmod{p}.$$

Now inserting (52)–(55) into (51), we obtain

$$(56) \quad \sum_{1 \leq i \leq k \leq p-1} \frac{(-1)^{k-1}}{ik} \binom{p-1}{k} \equiv -\frac{1}{2} H_{p-1,2} + p \sum_{k=1}^{p-1} \frac{H_k^2}{k}.$$

The equality (50) and the congruence (56) give

$$(57) \quad \sum_{k=1}^{p-1} \frac{H_k^2}{k} \equiv \frac{3}{2p} \sum_{k=1}^{p-1} \frac{1}{k^2} \pmod{p^2}.$$

The congruences (57) and (48) of Lemma 2.10 yield

$$(58) \quad \sum_{k=1}^{p-1} \frac{H_k^2}{k} \equiv -\frac{3}{p^2} \sum_{k=1}^{p-1} \frac{1}{k} \pmod{p^2}.$$

Finally, the congruences (57) and (58) complete proof of Theorem 1.1. $\square$

Remark. From the identity

$$H_{k-1}^3 = \left(H_k - \frac{1}{k} \right)^3 = H_k^3 - 3 \frac{H_k^2}{k} + 3 \frac{H_k}{k^2} - \frac{1}{k^3}$$

immediately follows that

$$\sum_{k=1}^{p-1} \frac{H_k^2}{k} - \sum_{k=1}^{p-1} \frac{H_k}{k^2} = \frac{1}{3} (H_{p-1}^3 - H_{p-1,3}).$$

Inserting in the right hand side of the above identity the congruences $H_{p-1} \equiv 0 \pmod{p^2}$ and $H_{p-1,3} \equiv -\frac{6p^2 B_{p-5}}{5} \pmod{p^3}$ from [9, Theorem 5.1(a) with $k = 3$], we find that for a prime $p \geq 7$,

$$\sum_{k=1}^{p-1} \frac{H_k^2}{k} - \sum_{k=1}^{p-1} \frac{H_k}{k^2} \equiv \frac{2p^2 B_{p-5}}{5} \pmod{p^3}.$$

However, the determination of $\sum_{k=1}^{p-1} \frac{H_k^2}{k} \pmod{p^3}$ it seems to be a difficult problem.

REFERENCES

- [1] E. Alkan, Variations on Wolstenholme's theorem, *Amer. Math. Monthly* **101** (1994), 1001–1004.
- [2] M. Bayat, A generalization of Wolstenholme's Theorem, *Amer. Math. Monthly* **104** (1997), 557–560.
- [3] A. Granville, Arithmetic properties of binomial coefficients. I. Binomial coefficients modulo prime powers, in *Organic Mathematics–Burnaby, BC 1995*, CMS Conf. Proc., vol. 20, American Mathematical Society, Providence, RI, 1997, 253–276.
- [4] V. Hernández, Solution IV of problem 10490 (a reciprocal summation identity), this *Amer. Math. Monthly* **106** (1999), 589–590.
- [5] R. Meštrović, A proof of a conjecture by Sun on congruence for harmonic numbers, preprint arXiv: 1108.1171v1 [math.NT] (2011).
- [6] R. Meštrović, On the mod p^7 determination of $\binom{2p-1}{p-1}$, preprint arXiv: 1108.1174v1 [math.NT] (2011).
- [7] J. Mináč, Newton's identities once again, *Amer. Math. Monthly* **110** (2003), 232–234.
- [8] H. Pan and Z. W. Sun, New identities involving Bernoulli and Euler polynomials, *J. Combin. Theory Ser. A* **113** (2006), 156–175.
- [9] Z. H. Sun, Congruences concerning Bernoulli numbers and Bernoulli polynomials, *Discrete Appl. Math.* **105** (2000), 193–223.
- [10] Z. W. Sun, Arithmetic theory of harmonic numbers, *Proc. Amer. Math. Soc.*, article in press, preprint arXiv: 0911.4433v3 [math.NT] (2009).
- [11] Z. W. Sun and R. Tauraso, New congruences for central binomial coefficients, *Adv. in Appl. Math.* **45** (2010), 125–148.
- [12] R. Tauraso, Problem 11382, *Amer. Math. Monthly* **115**, August–September, (2008).
- [13] R. Tauraso, New harmonic number identities with applications, *Séminaire Lotharingien de Combinatoire* **63** (2010), Article B63g.
- [14] R. Tauraso, More congruences for central binomial coefficients, *J. Number Theory* **130** (2010), 2639–2649.
- [15] D. B. Tyler, Solution of problem 11382, *Amer. Math. Monthly* **118** (2011), 85–86.
- [16] J. Zhao, Wolstenholme type theorem for multiple harmonic sums, *Int. J. Number Theory* **4** (2008), 73–106.
- [17] X. Zhou and T. Cai, A generalization of a curious congruence on harmonic sums, *Proc. Amer. Math. Soc.* **135** (2007), 1329–1333.

DEPARTMENT OF MATHEMATICS, MARITIME FACULTY, UNIVERSITY OF MONTENEGRO, DOBROTA 36, 85330 KOTOR, MONTENEGRO ROMEO@AC.ME